

AD_____

Award Number: W81XWH-09-1-0289

TITLE: Circadian Biology and Sleep: Missing Links in Obesity and Metabolism

PRINCIPAL INVESTIGATOR: Jeffrey M. GIMBLE, M.D., Ph.D.

CONTRACTING ORGANIZATION: Louisiana State University System
Baton Rouge, LA 70808

REPORT DATE: May 2009

TYPE OF REPORT: Final Proceeding

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE 29-05-2009		2. REPORT TYPE Final Proceedings		3. DATES COVERED 26 APR 2009 - 29 APR 2009	
4. TITLE AND SUBTITLE Circadian Biology and Sleep: Missing Links in Obesity and Metabolism				5a. CONTRACT NUMBER W81XWH-09-1-0289	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Jeffrey M. GIMBLE, M.D., Ph.D. Email: i k o d r g l o B r d t e Q f w				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Louisiana State University System Baton Rouge, LA 70808				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT The 2009 Pennington Scientific Symposium entitled "Circadian Biology and Sleep": Missing Links in Obesity and Metabolism? Was held at the Lod Cook Conference Center on the Louisiana State University Campus on 4/27-28, 2009. Over 25 invited speakers and their guests together with ~15 to 20 faculty members from the Pennington Biomedical Research Center attended the sessions which integrated current findings in the fields of circadian biology, metabolism, nutrition, and sleep. In addition to the keynote address by Dr. Joseph Takahashi, a total of 14 full talks and 2 short talks were presented. A round-table discussion was conducted at the conclusion of each day's meeting. Each speaker was given the option to submit his/her talk in manuscript format to the journal <i>Obesity Reviews</i> ; over 75% of the speakers have indicated that they will publish in this special supplement issue.					
15. SUBJECT TERMS Circadian biology; metabolism; molecular biology; obesity; sleep					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 42	19a. NAME OF RESPONSIBLE PERSON USAMRMC
a. REPORT U	b. ABSTRACT U	c. THIS PAGE U			19b. TELEPHONE NUMBER (include area code)

Introduction:

We now live in a mechanized world that is increasingly divorced from the physical environment. With artificial lighting, we no longer experience a 10-12 hour period of true darkness at night. Our sleep cycles and activity periods have become divorced from the earth's rotational rhythm. Night shift workers might be eating lunch at 2 AM and sleeping at 2 PM. With modern air travel, we now experience "jet lag" as our bodies are forced abruptly to acclimate to new time zones and altered light:dark cycles. There will be increasing circadian stresses placed on the U.S. civilian population's lifestyle as globalization forces factories, retail commerce, and computer-based jobs towards a "24/7" operating schedule. Nevertheless, the comprehensive integration of the fields of circadian biology, sleep, metabolism, nutrition, and human physiology remains a challenge for investigators, clinicians, and administrators in both the civilian and military settings.

Body:

To address the important questions concerning the role of circadian mechanisms in human disease and to foster inter-disciplinary communication and collaboration, a symposium entitled "*Circadian Biology and Sleep: Missing Links in Obesity and Metabolism?*" was held at the Lod Cook Conference Center on the campus of the Louisiana State University (Baton Rouge, LA) between April 26-29, 2009. The co-Chairs for the meeting were Andrew Young, PhD, from the U.S. Army Research Institute of Environmental Medicine, Molly Bray, PhD, from the Baylor College of Medicine, and Jeffrey M. Gimble, MD PhD, from the Pennington Biomedical Research Center. Each of these investigators has expertise in the field of human performance, circadian biology, adipose tissue, and/or obesity research.

The following distinguished panel of scientists from the fields of circadian biology, sleep, human performance, and obesity research attended this meeting and presented their talks entitled as listed below.

Keynote Address: Joseph Takahashi, PhD (Northwestern University):
Uncovering the Molecular Basis of Circadian Biology

Full Talks:**Joseph Bass, MD, PhD**

Northwestern University

Obesity and Metabolism in the Clock Mutant Mice: Lessons for Human Metabolism

Molly Bray, PhD

Baylor College of Medicine

Genetic Variation in Circadian Genes: Relation to Obesity and Adipocyte Biology

Andrew Butler, PhD

The Scripps Research Institute

The role of melanocortin neuronal pathways in circadian biology: a new homeostatic output involving melanocortin-3 receptors?

Nicolas Cermakian, PhD

McGill University

Regulation of central and peripheral circadian rhythms in mice and humans

James Gangwisch, PhD

Columbia University

The Epidemiological Relationships Between Sleep Patterns and Metabolic Disorders

Jeff Gimble, MD, PhD

Pennington Biomedical Research Center

Food Entrainment and the Circadian Regulatory Apparatus in Peripheral Tissues

Carla Green, PhD

University of Virginia

Nocturnin: A Novel Circadian and Metabolic Regulator

Jonathan Johnston, PhD

University of Surrey

Adipocyte rhythmicity in vitro

Tracy Rupp, PhD

Walter Reed Army Institute of Research

Long-term effects of nightly sleep duration on cognitive performance, alertness, metabolism and obesity

Jared Rutter, PhD

University of Utah

Nutrient sensing and signaling: Roles in energy balance and elsewhere

Albert Stunkard, MD

University of Pennsylvania

A Biobehavioral Model of the Night Eating Syndrome

Gregory Sutton, PhD

Pennington Biomedical Research Center

Maternal Imprinting and Metabolism: A Circadian Relationship?

Alexandros Vgontzas, MD

Penn State College of Medicine

Meaningful subtyping of Obesity based on Sleep Disturbances

Martin Young, PhD

Baylor College of Medicine

The Circadian Biology of Cardiac Tissues: A Metabolic Model for Understanding Human Metabolic Syndrome

Short Talks:

Karyn Esser, PhD

University of Kentucky,

Circadian Biology of Skeletal Muscle

Alok Gupta, MD

Pennington Biomedical Research Center,
Circadian Variation of Blood Pressure in Obese and Diabetic Subjects

Each of the invited speakers was encouraged to bring a junior colleague or collaborator to the meeting to participate in the proceedings and a total of 11 additional individuals from academia and industry attended the symposium. Additionally, the Pennington Biomedical Research Center faculty were invited to participate in the conference. Between 10-20 of the full time faculty members at the Center joined the meeting in addition to post-doctoral fellows, research associates, and staff members. The meeting sponsors had invited Dr. Michael Twery, a sleep specialist and program officer at the National Heart, Lung, and Blood Institute in Bethesda, MD, to attend. Although he had originally accepted and was on the program, his commitments at the NIH dealing with the stimulus package forced him to change his plans. Likewise, Dr. Andrew Young's other commitments prevented him from attending. In his place, Thomas Balkin, PhD, Chief, Department of Behavioral Biology, Walter Reed Army Institute for Research, attended the meeting and chaired one of the symposium sessions.

Arne Astrup, the editor in chief of the prestigious journal ***Obesity Reviews***, agreed to publish papers from the symposium. Each of the presenting scientists was invited to contribute an article based on their presentation. These articles will appear in a special issue of the journal which has an ISI impact factor of 7.821, higher than any other publication in the specialized field of obesity research. These publications, along with an overview of the meeting prepared by the organizers, will serve to disseminate the findings and conclusions of the conference to the broader scientific community. At least 75% of the speakers have agreed to submit articles based on their presentations. The meeting co-chairs and Dr. Balkin will prepare a summary statement on the meeting objectives and outcomes for this special supplement to ***Obesity Reviews***.

Key Research Accomplishments:

1. State of the art review of:
 - a. Circadian Biology at the Basic and Translational Levels
 - b. Nutrition and Obesity
 - c. Sleep Medicine
2. Cross Training of Investigators From Different Disciplines
3. Fostering of Inter-Disciplinary Collaborations
4. Development of a Special Supplement Issue of *Obesity Reviews* with Manuscripts from the Symposium

Reportable Outcomes:


The following reportable outcomes have or will result from this funding:

Abstract Booklet (attached)

Obesity Review Special Supplement with manuscripts from >75% of the symposium speakers and summary statement from symposium co-chairs (to be completed)

Conclusions:

The symposium sponsored by the US Army Medical and Materiel Command offered an outstanding educational forum that allowed exchange of ideas between scientists from different disciplines coming from laboratories in the US, Canada, and the United Kingdom. The event highlighted the existing knowledge and future directions for circadian biology in the context of human disease and performance relevant to both the military and civilian population. In addition, the event initiated dialogues with the potential to lead to collaborative efforts between US Army and academic laboratories, both national and international. Much of the information related at the symposium will be made available to the scientific community through its publication in a Special Supplement Issue of *Obesity Reviews*. This opportunity was only possible through the generous support of the US Army.



Pennington Scientific Symposium

Circadian Biology and Sleep: Missing Links in Obesity and Metabolism?

April 27-28, 2009

Symposium Chairs

Molly Bray, PhD

Associate Professor of Pediatrics
Baylor College of Medicine
Children's Nutrition Research Center

Jeffrey Gimble, MD, PhD

Professor, Stem Cell Biology
Pennington Biomedical Research Center
Louisiana State University System

Andrew Young, PhD

Chief, Military Nutrition Division
U.S. Army Research Institute of Environmental Medicine

This symposium was coordinated by the
Division of Education

Pennington Biomedical Research Center
Louisiana State University System
6400 Perkins Road
Baton Rouge, LA 70808-4124
Phone: 225/763-2629 Fax: 225/763-3045

Phillip J. Brantley, PhD, Director
Anne Schulte, Assistant Director

Monday, April 27

SESSION CHAIR: Molly Bray, PhD - Molecular Mechanisms in Circadian Biology and Metabolism

8:00-8:30 **Claude Bouchard, PhD**
Pennington Biomedical Research Center: Mission, Structure, Programs and Financing

8:30-9:30 **Joseph Takahashi, PhD**
Northwestern University
Keynote Address: Uncovering the Molecular Basis of Circadian Biology

9:30-10:15 **Joseph Bass, MD, PhD**
Northwestern University Feinberg School of Medicine
Obesity and Metabolism in the Clock Mutant Mice: Lessons for Human Metabolism

10:15-10:30 **BREAK**

10:30-11:15 **Carla Green, PhD**
University of Virginia
Nocturnin: A Novel Circadian and Metabolic Regulator

11:15-12:00 **Jared Rutter, PhD**
University of Utah School of Medicine
Nutrient sensing and signaling: Roles in energy balance and elsewhere

12:00 – 1:00 **LUNCH**

SESSION CHAIR: Jeff Gimble, MD, PhD - Circadian Biology of Sleep

1:00-1:45 **Albert Stunkard, MD**
University of Pennsylvania
A Biobehavioral Model of the Night Eating Syndrome

1:45-2:30 **Nicolas Cermakian, PhD**
McGill University
Regulation of central and peripheral circadian rhythms in mice and humans

2:30-2:45 **BREAK**

2:45-3:30 **Jonathan Johnston, PhD**
University of Surrey
Adipocyte rhythmicity in vitro

3:30-4:15 **James Gangwisch, PhD**
Columbia University
The Epidemiological Relationships Between Sleep Patterns and Metabolic Disorders

4:15-5:00 **Round Table Discussion**

5:00 **Hospitality Suite**

6:30 **Tour and Dinner for Out of Town Attendees - PBRC**

9:00 **Hospitality Suite**

Tuesday, April 28

SESSION CHAIR: Tom Balkin, PhD – Central & Peripheral Metabolic Tissues and Circadian Mechanisms

8:30-9:15

Andrew Butler, PhD

The Scripps Research Institute

The role of melanocortin neuronal pathways in circadian biology: a new homeostatic output involving melanocortin-3 receptors?

9:15-10:00

Molly Bray, PhD

Baylor College of Medicine

Genetic Variation in Circadian Genes: Relation to Obesity and Adipocyte Biology

10:00 – 10:15

BREAK

10:15-11:00

Gregory Sutton, PhD

Pennington Biomedical Research Center. LSU System

Maternal Imprinting and Metabolism: A Circadian Relationship?

11:00-11:45

Jeff Gimble, MD, PhD

Pennington Biomedical Research Center, LSU System

Food Entrainment and the Circadian Regulatory Apparatus in Peripheral Tissues

11:45-12:45

Lunch

SESSION CHAIR: Michael Twery, PhD - Integrating Sleep, Metabolism, and Circadian Pathways

12:45-1:30

Martin Young, PhD

Baylor College of Medicine

The Circadian Biology of Cardiac Tissues: A Metabolic Model for Understanding Human Metabolic Syndrome

1:30-2:15

Tracy Rupp, PhD

Walter Reed Army Institute of Research

Long-term effects of nightly sleep duration on cognitive performance, alertness, metabolism and obesity

2:15-2:30

BREAK

2:30-3:15

Alexandros Vgontzas, MD

Penn State College of Medicine

Meaningful subtyping of Obesity based on Sleep Disturbances

3:15-3:45

Michael Twery, PhD

National Heart, Lung, and Blood Institute

The NIH Perspective

3:45 - 5:00

Consensus Discussion

Address pre-symposium “critical questions in the field” and any additional questions raised during the meeting. Develop consensus for meeting report document.

5:00

Hospitality Suite

7:00

Dinner - Lod Cook Alumni Center

9:00

Hospitality Suite

Uncovering the Molecular Basis of Circadian Biology

Joseph S. Takahashi, PhD

Howard Hughes Medical Institute, UT Southwestern Medical Center, Dallas, TX

Circadian rhythms represent an evolutionarily conserved adaptation to the environment that can be traced back to the earliest life forms. In animals circadian behavior can be analyzed as an integrated system - beginning with genes leading ultimately to behavioral outputs. In the last decade, the molecular mechanism of circadian clocks has been uncovered by the use of phenotype-driven (forward) genetic analysis in a number of model systems. Circadian oscillations are generated by a set of genes forming a transcriptional autoregulatory feedback loop. In mammals, these include: *Clock*, *Bmal1*, *Per1*, *Per2*, *Cry1*, *Cry2* and *Casein kinase 1 epsilon*. Another dozen candidate genes have been identified and play additional roles in the circadian gene network such as the feedback loop involving *Rev-erba*. Despite this remarkable progress, it is clear that a significant number of genes that strongly influence and regulate circadian rhythms in mammals remain to be discovered and identified. As part of a large-scale N-ethyl-N-nitrosourea (ENU) mutagenesis screen using a wide range of nervous system and behavioral phenotypes, we have identified new circadian mutants in mice. These include new alleles of known circadian genes as well as novel circadian loci such *Fbx/3*, an orphan member of the F-box protein family, which we have found to interact selectively with the CRY proteins to target them for degradation through the proteasome pathway.

The discovery of 'clock genes' also led to the realization that the capacity for circadian gene expression is widespread throughout the body. Using circadian gene reporter methods, one can demonstrate that most peripheral organs and tissues can express circadian oscillations in isolation, yet still receive and may require input from the dominant circadian pacemaker in the suprachiasmatic nucleus (SCN) *in vivo*. We have used tissue-specific, conditional gene expression methods to analyze the relative contributions of central and peripheral circadian oscillators to circadian organization. The cellular autonomy of circadian clocks has raised a number of questions concerning synchronization and coherence of rhythms at the cellular level as well as circadian organization at the systems level. I will discuss recent work that addresses these issues and that examines a number of levels of complexity within the circadian system.

References:

1. Lowrey, P.L. and J.S. Takahashi. 2004. Mammalian circadian biology: Elucidating genome-wide levels of temporal organization. *Annual Review of Genomics and Human Genetics* 5:407-441.
2. Takahashi, J.S., H.-K. Hong, C.H. Ko and E.L. McDearmon. 2008. The genetics of mammalian circadian order and disorder: Implications for physiology and disease. *Nature Reviews Genetics* 9:764-775
3. Takahashi, J.S. 2004. Finding new clock components: Past and future. *Journal of Biological Rhythms* 19:339-347.
4. Vitaterna, M.H., L.H. Pinto and J.S. Takahashi. 2006. Large-scale mutagenesis and phenotypic screens for the nervous system and behavior in mice. *Trends in Neuroscience* 29:233-240.
5. Panda, S., M.P. Antoch, B.H. Miller, A.I. Su, A.B. Schook, M. Straume, P.G. Schultz, S.A. Kay, J.S. Takahashi and J.B. Hogenesch. 2002. Coordinated transcription of key pathways in the mouse by the circadian clock. *Cell* 109:307-320.
6. Yoo, S.-H., S. Yamazaki, P.L. Lowrey, K. Shimomura, C.H. Ko, E.D. Buhr, S.M. Siepk, H.-K. Hong, W.-J. Oh, O.-J. Yoo, M. Menaker and J.S. Takahashi. 2004. PERIOD2::LUCIFERASE real-time reporting of circadian dynamics reveals persistent circadian oscillations in mouse peripheral tissues. *Proc. Natl. Acad. Sci. USA* 101:5339-5346.
7. Welsh, D.K., S.-H. Yoo, A.C. Liu, J.S. Takahashi and S.A. Kay. 2004. Bioluminescence imaging of individual fibroblasts reveals persistent, independently phased circadian rhythms of clock gene expression. *Current Biology* 14:2289-2295.

8. Siepka, S.M., S.-H. Yoo, J. Park, W. Song, V. Kumar, Y. Hu, C. Lee and J.S. Takahashi. 2007. Circadian mutant *Overtime* reveals F-box protein FBXL3 regulation of *Cryptochrome* and *Period* gene expression. *Cell* 129:1011-1023.
9. Liu, A.C. D.K. Welsh, C.H. Ko, H.G. Tran, E.E. Zhang, A.A. Priest, E.D. Buhr, O. Singer, K. Meeker, I.M. Verma, F.J. Doyle, J.S. Takahashi and S.A. Kay. 2007. Intercellular coupling confers robustness against mutations in the SCN circadian clock network. *Cell* 129:605-616.
10. Hong, H.-K., J.L. Chong, W. Song, E.J. Song, A.A. Jyawook, A.C. Schook, C.H. Ko and J.S. Takahashi. 2007. Inducible and reversible *Clock* gene expression in brain using the tTA system for the study of circadian behavior. *PLoS Genetics* 3:324-338.
11. McDearmon, E.L., K.N. Patel, C.H. Ko, J.A. Walisser, A.C. Schook, J.L. Chong, L.D. Wilsbacher, E-J Song, H-K Hong, C.A. Bradfield and J.S. Takahashi. 2006. Dissecting the functions of the mammalian clock protein BMAL1 by tissue-specific rescue in mice. *Science* 314:1304-1308.

Obesity and Metabolism in the Clock Mutant Mice: Lessons for Human Metabolism

Joseph T. Bass, MD, PhD
Northwestern University, Evanston, IL

The circadian system orchestrates the temporal organization of many aspects of physiology and metabolism in synchrony with the 24 hr rotation of the Earth. Like the metabolic system, the circadian system is a complex feedback network that involves interactions between the brain and peripheral tissues. While the core clock resides in the master pacemaker neurons of the suprachiasmatic (SCN) nucleus, the entire clock network is expressed within both extra-SCN regions of the brain and in peripheral tissues.

Emerging evidence suggests that circadian regulation is intimately linked to metabolic homeostasis and that circadian dysregulation contributes to obesity and diabetes. At the molecular level, we have recently found that rhythmic NAD⁺ biosynthesis completes a novel metabolic feedback loop through which the positive limb of the clock leads to activation of the NAD⁺-dependent deacetylase Sirt1 which in turn inhibits Per2 transcription. These results pinpoint NAD⁺ biosynthesis and Sirt1 activity as a critical node in the molecular integration of circadian and metabolic networks. Studies of the crosstalk between circadian and metabolic systems have opened new understanding of obesity and diabetes mellitus.

Nocturnin: A Novel Circadian and Metabolic Regulator

Carla Green, PhD
University of Virginia, Charlottesville, VA

Nocturnin is a deadenylase that controls mRNA expression in a circadian manner by degrading the poly-A tails of target mRNAs, leading to mRNA turnover or translational silencing (1, 2). Previously we reported that a mouse lacking Nocturnin was resistant to diet-induced obesity and hepatic steatosis (3). The lean phenotype was not due to increased activity, decreased food intake or a higher metabolic rate. Transcript analysis in liver showed alterations in genes associated with lipid uptake and utilization.

In an effort to discern the mechanism behind this lean phenotype, we examined digestive tract function. Our initial studies indicated that the Nocturnin KO mice a faster transit time for lipid, but not water, in a gut motility assay. We therefore hypothesized that Nocturnin has a role in the absorption of dietary lipid in bowel. In order to investigate the mechanism further, we exposed WT and KO mice to various dietary challenges. When subjected to either a ketogenic diet or food restriction, our KO mice lost more weight than their WT counterparts. When on a standard mouse chow, the KO mouse exhibited lower circulating beta-hydroxybutyrate – a finding consistent with altered lipid availability in the KO. Moreover, this latter discrepancy in the KO is not due to hepatocyte malfunction as hepatocyte analysis showed normal rates of both lipid uptake and beta-oxidation.

We then examined aspects of digestive tract function directly by giving the mice a lipid bolus and performing histological examinations. The lipid droplets formed along the jejunum of the small intestine were significantly larger in the Nocturnin KO mice than in the WT mice. Through a subsequent series of *in vivo* and *in vitro* studies, we demonstrated that the KO mice are deficient in their ability to take up lipids. These animals have significantly disrupted lipid trafficking in the enterocytes, resulting in decreased absorption via apoB-containing non-HDL lipoproteins. We propose that Nocturnin has a role in the absorption of dietary lipid in bowel, presumably by altering genes necessary for metabolism or digestion through circadian post-transcriptional modifications of targeted transcripts.

References:

1. Baggs, J. E., and C. B. Green. 2003. Nocturnin, a Deadenylase in *Xenopus laevis* Retina. A Mechanism for Posttranscriptional Control of Circadian-Related mRNA. *Curr Biol* 13:189-98.
2. Garbarino-Pico, E., S. Niu, M. D. Rollag, C. A. Strayer, J. C. Besharse, and C. B. Green. 2007. Immediate early response of the circadian polyA ribonuclease nocturnin to two extracellular stimuli. *RNA* 13:745-55.
3. Green, C. B., N. Douris, S. Kojima, C. A. Strayer, J. Fogerty, D. Lourim, S. R. Keller, and J. C. Besharse. 2007. Loss of Nocturnin, a circadian deadenylase, confers resistance to hepatic steatosis and diet-induced obesity. *Proc Natl Acad Sci U S A* 104:9888-93.

Nutrient Sensing and Signaling: Roles in Energy Balance and Elsewhere

Jared Rutter, PhD
University of Utah, Salt Lake City, UT

The PAS domain is a sequence motif originally identified in and named for the transcription factors Period, the Aryl hydrocarbon receptor (AHR), and SIM. The aryl hydrocarbon receptor is present in the cytoplasm of normal cells. Xenobiotics, such as dioxin, bind to the PAS domain of AHR, and cause its translocation to the nucleus and activation of specific target genes. The PAS domain has subsequently been found in a large number of proteins from all phyla. Many of these proteins are the sensor proteins of bacterial two-component systems, which are responsible for responding to many metabolic and environmental stimuli, such as oxygen tension, redox, and light. In general, the PAS domain is thought to be a regulator of the activity of other domains within the protein. My work over the last decade and the work of my lab over the last five years has been focused on the PAS domain and a few of the proteins that contain and are regulated by PAS domains.

NPAS2 and Clock are basic helix-loop-helix PAS transcription factors that function in the circadian clock (Antoch et al., 1997; King et al., 1997; Reick et al., 2001). Since the identification of NPAS2 by colleagues in the McKnight lab (Zhou et al., 1997), we hypothesized, by analogy with the aryl hydrocarbon receptor, that it would be a ligand-activated transcription factor. Toward identifying putative ligands, we expressed and purified the PAS domain-containing portion of the protein. Interestingly, the protein when expressed in bacteria co-purified with a chromophore with an absorbance maximum of 420nm. We identified the chromophore as heme and mapped the interaction to the PAS domain of NPAS2. We were able to show that carbon monoxide regulated DNA binding activity of NPAS2 and Clock through interaction with the bound heme moiety (Dioum et al., 2002). In the intervening seven years, others have repeated and extended on these studies (Ishida et al., 2008; Koudo et al., 2005; Mukaiyama et al., 2006; Uchida et al., 2005). It is quite clear that the in vitro observations are reproducible. What is much less clear is the physiological relevance of these observations in the control of circadian rhythms, although recent mutagenesis studies have implicated heme binding in transcriptional activity in cultured cells (Ishida et al., 2008).

PASK is an evolutionarily conserved nutrient-responsive kinase. PASK consists of a PAS regulatory domain followed by a canonical ser/thr kinase domain (Rutter et al., 2001). As with many other PAS domain proteins, the activity of PASK is controlled by metabolic status. Studies in yeast and mammalian cells have demonstrated that PASK activity is nutrient-regulated and controls the decision whether to store or utilize available nutrients (da Silva Xavier et al., 2004; Grose et al., 2007) (Smith and Rutter, 2007) (Rutter et al., 2002a). In order to understand the functions of PASK in mammals, we are studying *PASK*^{-/-} mice. *PASK*^{-/-} mice developed normally, but had altered metabolic parameters. The most dramatic observation was that *PASK*^{-/-} mice were resistant to high-fat diet induced obesity, hepatic steatosis and insulin resistance (Hao et al., 2007). We have focused on the role of PASK in regulating hepatic lipid metabolism and have identified a transcriptional pathway by which it might be mediating its effects. These findings suggest an important physiological role of PASK in regulating metabolism and controlling energy balance in mammals (Hao and Rutter, 2008). They also raise the possibility that PASK inhibition would be an effective therapeutic strategy for metabolic disease.

Previous colleagues and I proposed in 2002 that circadian rhythms might be, ancestrally and fundamentally, a metabolic cycle that has been reinforced by an elegant regulatory system of transcriptional, phosphorylation and protein degradation control (Rutter et al., 2002b). The connections between circadian rhythms and metabolism are certainly abundant and powerful (see (Green et al., 2008) for a recent review). As we have begun to understand some of the metabolic bases of obesity and diabetes, the issues of energetic efficiency, rhythmic activity of mitochondria and metabolic enzymes, rhythmic availability of energetic substrate and others have become very interesting areas where circadian rhythms might be interacting with metabolic regulation in highly disease-relevant scenarios.

References:

- Antoch, M.P., Song, E.J., Chang, A.M., Vitaterna, M.H., Zhao, Y., Wilsbacher, L.D., Sangoram, A.M., King, D.P., Pinto, L.H., and Takahashi, J.S. (1997). Functional identification of the mouse circadian Clock gene by transgenic BAC rescue. *Cell* 89, 655-667.
- da Silva Xavier, G., Rutter, J., and Rutter, G.A. (2004). Involvement of Per-Arnt-Sim (PAS) kinase in the stimulation of preproinsulin and pancreatic duodenum homeobox 1 gene expression by glucose. *Proc Natl Acad Sci U S A* 101, 8319-8324. Epub 2004 May 8317.
- Dioum, E.M., Rutter, J., Tuckerman, J.R., Gonzalez, G., Gilles-Gonzalez, M.A., and McKnight, S.L. (2002). NPAS2: a gas-responsive transcription factor. *Science* 298, 2385-2387.
- Green, C.B., Takahashi, J.S., and Bass, J. (2008). The meter of metabolism. *Cell* 134, 728-742.
- Grose, J.H., Smith, T.L., Sabic, H., and Rutter, J. (2007). Yeast PAS kinase coordinates glucose partitioning in response to metabolic and cell integrity signaling. *Embo J* 26, 4824-4830.
- Hao, H.X., Cardon, C.M., Swiatek, W., Cooksey, R.C., Smith, T.L., Wilde, J., Boudina, S., Abel, E.D., McClain, D.A., and Rutter, J. (2007). PAS kinase is required for normal cellular energy balance. *Proc Natl Acad Sci U S A* 104, 15466-15471.
- Hao, H.X., and Rutter, J. (2008). The role of PAS kinase in regulating energy metabolism. *IUBMB Life* 60, 204-209.
- Ishida, M., Ueha, T., and Sagami, I. (2008). Effects of mutations in the heme domain on the transcriptional activity and DNA-binding activity of NPAS2. *Biochem Biophys Res Commun* 368, 292-297.
- King, D.P., Zhao, Y., Sangoram, A.M., Wilsbacher, L.D., Tanaka, M., Antoch, M.P., Steeves, T.D., Vitaterna, M.H., Kornhauser, J.M., Lowrey, P.L., *et al.* (1997). Positional cloning of the mouse circadian clock gene. *Cell* 89, 641-653.
- Koudo, R., Kurokawa, H., Sato, E., Igarashi, J., Uchida, T., Sagami, I., Kitagawa, T., and Shimizu, T. (2005). Spectroscopic characterization of the isolated heme-bound PAS-B domain of neuronal PAS domain protein 2 associated with circadian rhythms. *Febs J* 272, 4153-4162.
- Mukaiyama, Y., Uchida, T., Sato, E., Sasaki, A., Sato, Y., Igarashi, J., Kurokawa, H., Sagami, I., Kitagawa, T., and Shimizu, T. (2006). Spectroscopic and DNA-binding characterization of the isolated heme-bound basic helix-loop-helix-PAS-A domain of neuronal PAS protein 2 (NPAS2), a transcription activator protein associated with circadian rhythms. *Febs J* 273, 2528-2539.
- Reick, M., Garcia, J.A., Dudley, C., and McKnight, S.L. (2001). NPAS2: An analog of clock operative in the mammalian forebrain. *Science* 293, 506-509.
- Rutter, J., Michnoff, C.H., Harper, S.M., Gardner, K.H., and McKnight, S.L. (2001). PAS kinase: An evolutionarily conserved PAS domain-regulated serine/threonine kinase. *Proc Natl Acad Sci USA* 98, 8991-8996.
- Rutter, J., Probst, B.L., and McKnight, S.L. (2002a). Coordinate regulation of sugar flux and translation by PAS kinase. *Cell* 111, 17-28.
- Rutter, J., Reick, M., and McKnight, S.L. (2002b). Metabolism and the Control of Circadian Rhythms. *Annual Review of Biochemistry* 71, 307-331.
- Smith, T.L., and Rutter, J. (2007). Regulation of Glucose Partitioning by PAS Kinase and Ugp1 Phosphorylation. *Mol Cell* 26, 491-499.

Uchida, T., Sato, E., Sato, A., Sagami, I., Shimizu, T., and Kitagawa, T. (2005). CO-dependent activity-controlling mechanism of heme-containing CO-sensor protein, neuronal PAS domain protein 2. *J Biol Chem* 280, 21358-21368.

Zhou, Y.D., Barnard, M., Tian, H., Li, X., Ring, H.Z., Francke, U., Shelton, J., Richardson, J., Russell, D.W., and McKnight, S.L. (1997). Molecular characterization of two mammalian bHLH-PAS domain proteins selectively expressed in the central nervous system. *Proc Natl Acad Sci USA* 94, 713-718.

A Biobehavioral Model of the Night Eating Syndrome

Albert J. Stunkard, MD
University of Pennsylvania, Pittsburgh, PA

The Night Eating Syndrome is an eating disorder consisting of evening hyperphagia (at least 25% of daily caloric intake after supper) and/or two awakenings at night with ingestions (1). It represents a delayed circadian rhythm of meal times of about 1.5 hours, with a similar delay in neuroendocrine functions (2). Prevalence of NES has varied considerably in the past depending upon the criteria which had changed. At the present time, the two criteria noted above have been widely accepted and data on prevalence is based on these criteria. In community samples it varies from 1.1% (3) to 1.6% (4) with higher prevalence reported among obese patients in weight loss samples (5) and even higher prevalence among preoperative bariatric surgery patients (5). One of the highest prevalences was found in psychiatric clinics with increased risk for patients with substance abuse disorders (6). The only population-based study of the prevalence of NES, based on the Swedish Twin Registry, found a prevalence of night eating among 4.6% of men and 3.4% of women (7). This, and other studies, found a strong relationship between body weight and the prevalence of night eating.

One large prospective study showed that nighttime overeaters gained significantly weight (6.2 kg) than did non-night eaters (1.7 kg; $p=0.03$) (8). NES is believed to result from an interaction of genetic vulnerability (7) and stress which has been consistently reported as a precipitating factor. The result of this interaction is an elevation in density of serotonin transporters in the midbrain, which transport serotonin from the synapse to the cell body, leaving a deficit of serotonin for neurotransmission (9). This deficit leads to the disruption of normal circadian rhythm of meal times and neuroendocrine functions. This model is supposed by four studies of selective serotonin reuptake inhibitors (SSRIs), which inhibit the reuptake of serotonin and increase its presence at the synapse (10-13). This treatment has been effective in controlling night eating. It also resulted in weight loss, the only situation in which SSRIs have produced weight loss and indicates that this action is via control of NES.

Despite the ease of diagnosis of NES and the presence of effective treatment, the disorder is not widely recognized. A survey of 103 night eaters indicated that 90% of them said that their doctors hadn't heard of NES and 40% said that their doctors had told them that their complaints were of no consequence (14).

References:

1. Allison KC, Lundgren JD, O'Reardon JP, Geliebter A, Gluck ME, Vinai P, Mitchell JE, Schenck CH, Howell MJ, Crow SJ, Engel S, Latzer Y, Tzischinsky O, Mahowald MW, Stunkard AJ. Proposed diagnostic criteria for night eating syndrome. *Int J Eat Disord*, In press.
2. Goel N, Stunkard AJ, Rogers NL, Van Dongen HPA, Allison KC, O'Reardon JP, Ahima RS, Cummings DE, Moonseong H, Dinges D. Circadian rhythm profiles in women with night eating syndrome. *J Biol Rhythms*. 2009;24:85.
3. Lamerz A, Kuepper-Nybelen J, Bruning N, Wehle C, Trost-Brinkhues G, Brenner H, Hebebrand J, Herpertz-Dahlmann B. Prevalence of obesity, binge eating and night eating in a cross-sectional field survey of 6-year-old children and their parents in a German urban population. *J Child Psychol Psychiatry*. 2005;46:385-393.
4. Striegel-Moore RH, Franko DL, Thompson D, Affenito S, Kraemer HC. Night Eating: Prevalence and demographic correlates. *Obes Res*. 2006;14:139-147.
5. Allison KC, Wadden TA, Sarwer DB, Fabricatore AN, Cserny CE, Gibbons LM, Stack RM, Stunkard AJ, Williams NN. Night eating syndrome and binge eating disorder among persons seeking bariatric surgery: prevalence and related features. *Obesity*. 2006;4:77S-82S.
6. Lundgren JD, Allison KC, Crow S, O'Reardon JP, Berg KC, Galbraith J, Martino NS, Stunkard AJ. Prevalence of the night eating syndrome in a psychiatric population. *Am J Psychiatry*. 2006;163:156-158.

7. Tholin S, Lindroos AK, Tynelius P, Akerstedt T, Stunkard AJ, Bulik C. Prevalence of night eating in obese and non-obese twins. *Obesity*. In press.
8. Gluck ME, Venti CA, Salbe AD, Krakoff J. Nighttime Eating: commonly observed and related to weight gain in an inpatient food intake study. *Am J Clin Nut*. 2008;88:900-905.
9. Frazer A. Pharmacology of antidepressants. *J Clin Psychopharmacol*. 1997;17(Suppl1):2S-18S.
10. Miyaoka T, Yasukawa R, Tsubouchi K, Miura S, Shimizu Y, Sukegawa T, Maeda T, Mizuno S, Kameda A, Uegaki J, Inagaki T, Horiguchi J. Successful treatment of nocturnal eating/drinking syndrome with selective serotonin reuptake inhibitors. *Int Clin Psychopharmacol*. 2003;18:175-177.
11. O'Reardon JP, Stunkard AJ, Allison KC. A clinical trial of sertraline in the treatment of Night Eating Syndrome. *Int J Eat Disord*. 2004;35:16-26.
12. Stunkard AJ, Allison KC, Lundgren JD, Martino NS, Heo M, Etemad B, O'Reardon JP. A paradigm for facilitating pharmacotherapy at a distance: sertraline treatment of the night eating syndrome. *J Clin Psychiatry*. 2006;67:1558-1572.
13. O'Reardon JP, Allison KC, Martino NS, Lundgren JD, Heo M, Stunkard AJ. A randomized placebo-controlled trial of sertraline in the treatment of the night eating syndrome. *Am J Psychiatry*. 2006;164:893-898.
14. Goncalves MD, Moore RH, Stunkard AJ, Allison KC. The treatment of night eating: the patient's perspective. In press.

Regulation of Central and Peripheral Circadian Rhythms in Mice and Humans

Cermakian, N.¹, James, F.O.¹, Fortier, E.E.^{1,2}, Dardente, H.¹, Rooney, J., Hardy, M.P.², Bélanger, V.¹, Charbonneau, S.¹, Villemain, A.¹, Labrecque, N.², Boivin, D.B.¹

¹ Douglas Mental Health University Institute, McGill University, Montreal, QC, Canada

² Maisonneuve-Rosemont Hospital Research Centre, University of Montreal, Montreal, QC, Canada

Many circadian rhythms are controlled by the central clock of the suprachiasmatic nucleus (SCN) of the hypothalamus¹. However, many other brain regions and most peripheral tissues have also been shown to function as autonomous clocks². The role of these peripheral clocks and their regulation by environmental and SCN-derived signals is still unclear. We address these questions in both mice and humans, and more specifically, we study circadian rhythms in cells of the immune system.

Our studies in mice focus on circadian rhythms in lymph nodes (LNs). T cell proliferation after antigen presentation in LNs is the first step in the adaptive immune response³. Serum levels of cytokines and circulating immune cells were previously shown to vary with a 24-hour circadian rhythm^{4,5}. Also, the persistence of rhythmic immune parameters under constant light conditions, under which no central clock-driven rhythms are present, suggested that there is a clock in the immune system⁵. We hypothesized that a circadian clock in LNs controls T cell function. Our aims were to identify clock gene expression in mouse LNs, investigate T cell proliferation rhythms, and examine LN rhythms in mice with a deficient clock. Adult male C57BL/6 mice were housed under a light:dark cycle, placed in constant darkness (DD), and sacrificed on the second day in DD every 4 h over 24 h. LNs were sampled and used to: i) Extract RNA and quantify clock gene expression by real-time PCR; or ii) Measure T cell proliferation following T cell receptor stimulation with an anti-CD3 antibody in vitro. Similar experiments were performed with *Clock* mutant mice and wild-type littermates. We show that clock genes are rhythmically expressed in lymph nodes and that the clock controls T cell proliferation. T cells undergo more divisions in response to stimulation when sampled from mice in the early night compared to other times of day, and this variation is lost in cells collected from *Clock* mutant mice. We are currently investigating intracellular signaling pathways downstream of the T cell receptor to assess the mechanisms behind this circadian rhythm in the proliferative response. This is the first evidence of a clock in LNs, and of control of the immune response by the molecular clockwork. Our results have uncovered a novel mode of regulation of T-cell proliferation. These studies will shed light on the links between circadian rhythms, immune response and cancer. They may also provide cues for more efficient vaccination strategies.

The rationale of our studies in humans is to use clock gene expression in peripheral blood mononuclear cells (PBMCs) as a model of peripheral oscillator, and to compare the regulation of central and peripheral rhythms under different environmental conditions. In a first study, we used an uninterrupted 72-h sampling period to compare the expression of peripheral circadian oscillators in PBMCs in the presence of a habitual sleep/wake cycle (LD) and under constant routine (CR) conditions⁶. Six healthy men (n=4) and women (n=2, follicular phase) aged 18-30 years (mean age \pm SEM: 23.7 \pm 1.6) maintained a regular 8-hour sleep episode for two weeks prior to the study. Repeated whole blood samples were drawn from an indwelling catheter during a 72-hour period including 40 h of LD where subjects were exposed to 118 \pm 8 lux of polychromatic light during daytime wake periods and slept in darkness, and a 32-h CR of sustained wakefulness and limited activity under dim light. Real-time PCR was used to assess expression of *PER1*, *PER2* and *BMAL1* in PBMCs isolated from whole blood samples every ~120 min, relative to *CDK4*. Plasma melatonin sampled every ~60 min and continuously (~1/min) sampled core body temperature (CBT) were used as markers of the central circadian pacemaker. The CBT minimum and the midpoint of melatonin concentration were well aligned to the sleep/wake cycle. *PER1* and *PER2* expression in PBMCs demonstrated significant circadian rhythmicity that peaked early after wake time and was comparable under LD and CR conditions. *BMAL1* expression was more variable, and peaked in the middle of the wake period under LD conditions and during the habitual sleep period under CR conditions. These results demonstrate that the pattern of clock gene expression in PBMCs can be evaluated over extended periods, and the pattern of *PER1* and *PER2* expression are in stable relationships with markers of the central circadian pacemaker.

Judicious light and darkness exposure throughout the day can promote the appropriate alignment of the endogenous hormonal rhythms to night shift work⁷. However, the synchronization of human peripheral

circadian oscillators to shifted sleep-wake schedules is currently unknown. Thus, building on the results of our previous experiment, we evaluated clock gene expression in PBMCs with respect to the simultaneous resetting of the plasma cortisol and melatonin rhythms throughout simulated night shift work⁸. Five healthy candidates (4 male, 1 female in follicular phase) aged (mean \pm SD) 24.9 \pm 4.8 years maintained stable sleep and meal schedules before the study start. Upon admission to the laboratory, sleep/wake schedules were delayed by 10 h to simulate nighttime “work”. The light intervention included exposure to polychromatic white light of (mean \pm SEM) 6,036 \pm 326 lux during 8-h night shifts and dim light exposure after each night shift with the use of sunglasses (5% visual light transmission). Phase of clock gene expression in PBMCs, and plasma melatonin and cortisol concentration were estimated from 24-h blood sampling periods performed before and after nine simulated night shifts. The expression of clock genes *PER1*, *PER2* and *BMAL1* was determined as above. Following nine days on the night schedule, melatonin and cortisol rhythms regained their conventional alignment with the shifted sleep/wake schedule. Changes in the pattern of clock gene expression were apparent when sampled as early as after three days on the shifted sleep/wake schedule. Following nine days on the night schedule, *PER1* and particularly *PER2* expression in PBMCs displayed significant circadian rhythmicity and were in a conventional alignment with the shifted sleep/wake schedule, with peak expression occurring during the wake period. This is the first demonstration of the entrainment of a human peripheral circadian oscillator to an atypical sleep-wake schedule. This line of investigation has important implications for understanding the medical disorders affecting night shift workers.

Funded by the Canadian Institutes of Health Research

References:

1. Cermakian, N. & Boivin, D. B. A molecular perspective of human circadian rhythm disorders. *Brain Res Brain Res Rev* 42, 204-220 (2003).
2. Schibler, U., Ripperger, J. & Brown, S. A. Peripheral circadian oscillators in mammals: time and food. *J Biol Rhythms* 18, 250-260 (2003).
3. Janeway, C. A., Travers, P., Walport, M. & Shlomchik, M. J. *Immunobiology: the immune system in health and disease*. 6th ed. edn, (Garland Science, 2005).
4. Born, J., Lange, T., Hansen, K., Molle, M. & Fehm, H. L. Effects of sleep and circadian rhythm on human circulating immune cells. *J Immunol* 158, 4454-4464 (1997).
5. Depres-Brummer, P., Bourin, P., Pages, N., Metzger, G. & Levi, F. Persistent T lymphocyte rhythms despite suppressed circadian clock outputs in rats. *Am J Physiol* 273, R1891-1899 (1997).
6. James, F. O., Boivin, D. B., Charbonneau, S., Belanger, V. & Cermakian, N. Expression of clock genes in human peripheral blood mononuclear cells throughout the sleep/wake and circadian cycles. *Chronobiol Int* 24, 1009-1034 (2007).
7. Boivin, D. B., Tremblay, G. M. & James, F. O. Working on atypical schedules. *Sleep Med* 8, 578-589, doi:S1389-9457(07)00115-3 (2007).
8. James, F. O., Cermakian, N. & Boivin, D. B. Circadian rhythms of melatonin, cortisol, and clock gene expression during simulated night shift work. *Sleep* 30, 1427-1436 (2007).

Adipocyte Rhythmicity In Vitro

Jonathan Johnston, PhD
University of Surrey, United Kingdom

White adipose tissue (WAT) plays a key role in multiple aspects of physiology. In addition to storing energy in the form of triglycerides, WAT is now known to act as an active endocrine tissue via the synthesis and secretion of adipokines (1). Many aspects of endocrine physiology and metabolism are under the control of endogenous circadian clocks and there is now growing evidence for the circadian regulation of adipose function (2-5).

Analysis of transgenic mouse models provides a compelling physiological link between molecular components of the circadian clock and metabolism. Even though the resulting phenotype appears partially dependent on genetic background, mutation of clock-related genes consistently results in a metabolic dysregulation, including dyslipidaemia (6-9). Similar genetic links to metabolism have been found in humans, with clock gene polymorphisms being associated with obesity, metabolic syndrome and type 2 diabetes mellitus (10-12).

Within WAT, rhythmic gene expression is widespread and an estimated 20% of the murine WAT transcriptome, including adipokine mRNA, exhibits diurnal fluctuation (2,13-15). Furthermore, circadian regulation of endocrine activity is likely to be an important output of WAT, as studies of both rodents and humans have revealed 24-hour variation of adipokine concentration in the blood (16-19). Despite this emerging understanding of adipose rhythmicity, a number of key issues remain unanswered. Of these, my group is particularly focussed on *in vitro* and human physiological models.

To date, most studies of adipose rhythmicity have analysed *ex vivo* WAT derived from rodents maintained on a light-dark cycle. WAT is heterogeneous and contains multiple cell types in addition to the primary endocrine cells, the adipocytes. *In vivo*, WAT is subject to multiple rhythmic input signals, which are both humoral and neuronal in nature (20-23). Finally, there is a possibility of environmental 'masking' of physiological rhythms when individuals are housed in fluctuating environmental conditions. The cellular basis of reported adipose rhythms is therefore unclear and in need of further investigation. A major focus of current work is to characterise rhythmicity in isolated pre-adipocyte and adipocyte cells. Many of these studies are being performed in collaboration with Prof Gary Frost and other colleagues at Imperial College, London.

There have to date only been limited attempts to translate rodent data on WAT rhythms to human studies. I am currently leading a study, in collaboration with Prof Debra Skene and Dr Denise Robertson, to evaluate endocrine rhythms and WAT gene expression in physiologically characterised human subjects.

I will present preliminary data from the above studies and provide an overview of the links between clocks, sleep, metabolism and nutrition within the Sleep and Chronobiology group at the University of Surrey.

Research funded by BBSRC, Diabetes-UK and Stockgrand UK

References:

1. Trujillo and Scherer 2006 *Endocrine Rev* 27:762-778
2. Bray and Young 2006 *Obesity Rev* 8:169-181
3. Hastings et al 2007 *J Endocrinol* 195:187-198
4. Ramsey et al 2007 *Annu Rev Nutr* 27:219-240
5. Zvonic et al 2007 *Obesity* 15:539-543
6. Turek et al 2005 *Science* 308:1043-1045

7. Oishi et al 2006 FEBS Lett 580:127-130
8. Kondratov et al 2006 Genes Dev 20:1868-1873
9. Green et al 2007 Proc Natl Acad Sci USA 104:9888-9893
10. Scott et al 2008 Int J Obes 32:658-62
11. Woon et al 2007 Proc Natl Acad Sci USA 104:14412-14417
12. Monteleone et al 2008 Neurosci Lett 435:30-33
13. Ptitsyn et al 2006 PLoS Comput Biol 2(3):e16
14. Zvonic et al 2006 Diabetes 55:962-970
15. Ando et al 2005 Endocrinology 146:5631-5636
16. Sinha et al 1996 J Clin Invest 97:1344-1347
17. Saad et al 1998 J Clin Endocrinol Metab 83:453-459
18. Kalsbeek et al 2001 Endocrinology 142:2677-2685
19. Gavrilu et al 2003 J Clin Endocrinol Metab 88:2838-2843
20. Balsalobre et al 2000 Science 289:2344-2347
21. Pévet et al 2006 Chronobiol Int 23:39-51
22. Bartness et al 2001 J Biol Rhythms 16:196-204
23. Kreier et al 2006 Endocrinology 147:1140-1147

The Epidemiological Relationship Between Sleep Patterns and Metabolic Disorders

James E. Gangwisch, Ph.D.

Columbia University, College of Physicians and Surgeons

Department of Psychiatry, Division of Cognitive Neuroscience, New York, NY

In this country we are experiencing parallel trends of increasing body mass index, increasing prevalence of hypertension and diabetes, and decreasing average sleep duration. There is reason to believe that these trends are linked by a mismatch between our modern lifestyles and our metabolic regulatory systems that evolved to be functional when the lifestyles of our hunter-gatherer ancestors were quite different from today. The thrifty genotype is hypothesized to have evolved through natural selection to be expressed during seasons of high food availability to induce high blood insulin, insulin resistance, and glucose-intolerance to facilitate increased fat deposition to adaptively prepare for seasons of food scarcity. The central biological clock evolved to synchronize activity, consumption, and rest to the circadian and circannual cycles using hormones and the autonomic nervous system. To generate and organize autonomic rhythms, the SCN requires repeated metabolic cues from light exposure, sleep, activity, and nutrient intake. Reliable environmental cues of the impending seasons for our nomadic ancestors would have been the length of the daily photoperiod that changed in a precise and predictable fashion throughout the year and the intake of seasonally available nutrients. Longer photoperiods in summer would have corresponded with shorter sleep durations and higher availability of food and simple sugars. Short sleep durations could therefore help trigger the expression of the thrifty genotype to increase caloric intake and fat deposition. Today we can have year round short sleep durations with electric lights and year round availability of highly palatable foods and simple sugars, so we can have year round fat accumulation. The seasonal expression of the thrifty genotype is therefore consistent with metabolic changes resulting from chronically short sleep durations leading to diseases associated with the metabolic syndrome such as obesity, diabetes, and hypertension.

Experimental studies have found sleep deprivation to increase appetite by decreasing leptin and increasing ghrelin levels. Inadequate sleep and poor-quality sleep are associated with obesity in children, adolescents, and adults studied in case-control, cross-sectional, and longitudinal studies¹⁻³. These relationships are stronger in children, adolescents and young adults. Associations in the elderly have been found with more accurate measures of sleep duration and adiposity.

Sleep deprivation in experimental studies has been shown to decrease glucose tolerance and compromise insulin sensitivity. It has been suggested that habitually short sleep durations could lead to insulin resistance by increasing sympathetic nervous system activity, raising evening cortisol levels, and decreasing cerebral glucose utilization. The increased burden on the pancreas from insulin resistance can, over time, compromise β -cell function and lead to type 2 diabetes. Longitudinal studies have found U-shaped relationships between sleep duration and diabetes incidence, with both short and long sleep durations being associated with diabetes incidence.⁴ It is likely that long sleep duration occurs in parallel to, and as a consequence of, diabetes and other conditions associated with chronic inflammation. Proinflammatory cytokines contribute toward sleepiness and fatigue, have deleterious effects on both glucose homeostasis and β -cell function, and are elevated in obesity and in conditions in which the primary pathogenic mechanism is insulin resistance.

Sleep restriction has been shown to acutely increase blood pressure and sympathetic nervous system activity in both normotensive and hypertensive subjects. Blood pressure dips by an average of 10 to 20% during sleep, so shorter sleep durations increase hemodynamic load by raising average 24-hour blood pressure and heart rate. Shorter sleep durations also result in longer exposures to waking physical and psychosocial stressors. Increased exposure to stress has been shown to promote salt appetite and suppress renal salt-fluid excretion. Long-term exposure to increased total 24-hour hemodynamic load associated with short sleep durations can lead to structural adaptations that gradually reset the entire cardiovascular system to operate at an elevated pressure equilibrium. Short sleep duration has been found to be associated with the prevalence of hypertension in cross-sectional studies and the incidence of hypertension in longitudinal studies.⁵ A u-shaped relationship between sleep duration and mortality has consistently been found in epidemiologic studies, with both short and long sleep durations being associated with increased mortality. The association between long sleep and mortality has been viewed as justification by some investigators to recommend sleep

restriction to decrease mortality risk despite the fact that no published laboratory or epidemiologic studies have demonstrated a possible mechanism identifying long sleep as a cause of either morbidity or mortality. In analyses stratified by age, the u-shaped relationship between sleep duration and mortality was found only in elderly subjects and not in middle-aged subjects.⁶ Results from these analyses indicate that the U-shaped relationship is heavily influenced by deaths in elderly subjects and by the measurement of sleep durations closely before death. Subjects' sleep durations reported closely before death are unlikely to be indicative of their sleep durations over the course of their lifetimes, when they develop chronic diseases that eventually contribute toward death. It is likely that the sleep durations of the elderly subjects who died were affected by the presence of medical conditions that eventually contributed toward their deaths. The majority of deaths in elderly subjects are attributable to major cardiovascular diseases, cancer, and diabetes. The inflammatory process has been shown to play key roles in both the pathogenesis and pathophysiology of cancer and of metabolic disorders such as cardiovascular disease and type 2 diabetes. The elderly subjects' immune responses to these conditions could have resulted in a predominance of pro-inflammatory cytokines that functioned to lengthen sleep, implying that long sleep duration is unlikely to contribute toward mortality but, rather, is a consequence of conditions associated with chronic inflammation and age-related sleep changes.

References:

1. Gangwisch JE, Malaspina D, Boden-Albala B, Heymsfield SB. Inadequate sleep as a risk factor for obesity: Analyses of the NHANES I. *SLEEP* 2005;28(10), 1265-1272.
2. Patel SR, Malhotra A, White DP, Gottlieb DJ, Hu FB. Association between reduced sleep and weight gain in women. *American Journal of Epidemiology* 2006;164:947-54.
3. Patel SR, Hu FB. Short sleep duration and weight gain: A systematic review. *Obesity* 2008;16:643-653.
4. Gangwisch JE, Heymsfield SB, Boden-Albala B, Buijs RM, Kreier F, Pickering TG, Rundle AG, Zammit GK, Malaspina D. Sleep duration as a risk factor for diabetes incidence in a large U.S. sample. *SLEEP* 2007;30(12),1667-1673.
5. Gangwisch JE, Heymsfield SB, Boden-Albala B, Buijs RM, Kreier F, Pickering TG, Rundle AG, Zammit GK, Malaspina D. Short sleep duration as a risk factor for hypertension: Analyses of the First National Health and Nutrition Examination Survey. *Hypertension* 2006;47(5),833-839.
6. Gangwisch JE, Heymsfield SB, Boden-Albala B, Buijs RM, Kreier F, Pickering TG, Rundle AG, Zammit GK, Malaspina D. Sleep duration associated with mortality in elderly, but not middle-aged adults in a large U.S. sample. *SLEEP* 2008;31(8),1087-1096.

The Role of Melanocortin Neuronal Pathways in Circadian Biology: A New Homeostatic Output Involved Melanocortin-3 Receptors?

Karima Begriche¹, Gregory M. Sutton², Jidong Fang³ and Andrew Butler¹, PhD

¹ Department of Metabolism and Aging, The Scripps Research Institute, Jupiter, FL

² Pennington Biomedical Research Center, LSU System, Baton Rouge, LA.

³ Department of Psychiatry, Pennsylvania State University College of Medicine

Energy homeostasis involves maintaining a balance of appetite with energy expenditure. The neural melanocortin system has a critical role in this process, and is involved in regulating both satiety and energy expenditure^{1,2}. In the brain, the melanocortin system is composed of neurons that secrete the endogenous melanocortin receptor ligands, and/or that express melanocortin receptors. The melanocortin receptor family is comprised of 5 g-protein coupled receptors (Mc1-5r), with Mc3r and Mc4r expressed in areas of the central nervous system involved in energy homeostasis. The melanocyte stimulating hormones (α -, β - and γ -MSH) exhibit agonist activity, and are products of the post-translation processing of proopiomelanocortin (Pomc). Agouti-related peptide (AgRP) was originally described as antagonist for the Mc3r and Mc4r, although this neuropeptide also acts as an inverse agonist. Neurons that express either *Pomc* or *AgRP* mRNA are located in the arcuate nucleus of the hypothalamus, and send projections to other neural centers involved in regulating complex behavior and the neuroendocrine and autonomic outputs that govern metabolism and energy expenditure. The hypothalamic melanocortin system responds rapidly to food intake in rats restricted to a once-daily feeding regime, suggesting acute regulation by peripheral inputs of meal intake³. The regulation of satiety and energy expenditure by melanocortin ligands predominantly involves Mc4r. Activation of Mc4r reduces food intake and increases energy expenditure^{1,2}. However, while having an important function in maintaining normal adiposity⁴, the exact role of the Mc3r in energy balance regulation has remained enigmatic.

My laboratory recently reported evidence suggesting that Mc3r are involved in entrainment to food intake⁵. The rationale for examining the role of melanocortins in food entrainment was based on the early studies demonstrating that hypothalamic lesions significantly impair entrainment of rats to a single meal⁶. Using *Mc3r*^{-/-} mice backcrossed >10 generations onto the C57BL/6J (B6) strain³, we observed significant impairment of entrainment to restricted feeding (RF: food available 1300-1700h in mice housed in a 12 h light:dark setting, with lights on 0600-1800h). Several readouts of activity and arousal were used to monitor entrainment. First, we observed using wheel cages that the development of food anticipatory activity (FAA) was significantly impaired in *Mc3r*^{-/-} mice. Second, we examined spontaneous locomotor activity (SLA) using beam breaks in the x- and z-axes using the Comprehensive Laboratory Animal Monitoring System (CLAMS). We observed a peak in SLA in control B6 mice subject to RF centered on food presentation which was markedly reduced in *Mc3r*^{-/-} mice. Using electroencephalography (EEG) and electromyography (EMG), we then observed that a period of wakefulness in control mice coinciding with FAA and anticipating meal presentation was absent in *Mc3r*^{-/-} mice.

FAA is dependent on the activity of genes encoding proteins involved in maintaining the rhythm of molecular clocks. Mice lacking either the Clock paralog neuronal PAS domain protein 2 (*Npas2*) or Period2 (*Per2*) exhibit impaired FAA when subject to restricted feeding protocols^{7,8}. We observed abnormal rhythms of expression of critical clock genes in the forebrain of *Mc3r*^{-/-} mice. Importantly, we were able to demonstrate similar abnormal rhythms of clock activity and impaired adaptation to RF in wild type B6 mice administered intracerebroventricularly with AgRP, a Mc3r antagonist/inverse agonist.

Collectively, these data suggest that one function of the Mc3r is to regulate inputs into systems involved in the development of behaviors that are entrained to food availability. We have also determined that Mc3r have a critical role in maintaining a normal rhythm of activity in the liver clock during RF. Using the same technology applied to the Mc4r⁹, we are now developing a condition mutant, the LoxTB-Mc3r mouse, to further define the role of neural Mc3r in regulating entrainment.

Acknowledgements: This work was supported by the Pennington Biomedical Research Foundation and the NIDDK (DK073189 to AB). The author thanks Dr Jeff Gimble (Pennington Biomedical Research Center, LSU System), Matthias Tschöp (Department of Psychiatry, Obesity Research Centre, University of Cincinnati) for their help with these experiments.

References:

1. Cone RD. Studies on the physiological functions of the melanocortin system. *Endocr Rev.* 2006 Dec;27(7):736-49.
2. Cone RD. Anatomy and regulation of the central melanocortin system. *Nat Neurosci.* 2005 May;8(5):571-8.
3. Sutton GM, Trevaskis JL, Hulver MW, McMillan RP, Markward NJ, Babin MJ, Meyer EA, Butler AA. Diet-genotype interactions in the development of the obese, insulin-resistant phenotype of C57BL/6J mice lacking melanocortin-3 or -4 receptors. *Endocrinology.* 2006 May;147(5):2183-96.
4. Johnstone LE, Fong TM, Leng G. Neuronal activation in the hypothalamus and brainstem during feeding in rats. *Cell Metab.* 2006 Oct;4(4):313-21.
5. Sutton GM, Perez-Tilve D, Nogueiras R, Fang J, Kim JK, Cone RD, Gimble JM, Tschöp MH, Butler AA. The melanocortin-3 receptor is required for entrainment to meal intake. *J Neurosci.* 2008 Nov 26;28(48):12946-55.
6. Choi S, Wong LS, Yamat C, Dallman MF. Hypothalamic ventromedial nuclei amplify circadian rhythms: do they contain a food-entrained endogenous oscillator? *J Neurosci.* 1998 May 15;18(10):3843-52.
7. Dudley CA, Erbel-Sieler C, Estill SJ, Reick M, Franken P, Pitts S, McKnight SL. Altered patterns of sleep and behavioral adaptability in NPAS2-deficient mice. *Science.* 2003 Jul 18;301(5631):379-83.
8. Feillet CA, Ripperger JA, Magnone MC, Dulloo A, Albrecht U, Challet E. Lack of food anticipation in *Per2* mutant mice. *Curr Biol.* 2006 Oct 24;16(20):2016-22.
9. Balthasar N, Dalgaard LT, Lee CE, Yu J, Funahashi H, Williams T, Ferreira M, Tang V, McGovern RA, Kenny CD, Christiansen LM, Edelstein E, Choi B, Boss O, Aschkenasi C, Zhang CY, Mountjoy K, Kishi T, Elmquist JK, Lowell BB. Divergence of melanocortin pathways in the control of food intake and energy expenditure. *Cell.* 2005 Nov 4;123(3):493-505.

Genetic Variation in Circadian Genes: Relation to Obesity and Adipocyte Biology

Molly Bray, PhD
Baylor College of Medicine, Houston, TX

Obesity is one of the most profound public health problems today, and though much has been learned regarding the regulation of body weight and the development of adiposity, the prevalence of obesity continues to rise. Simplistic explanations based on nutritional over-consumption and/or poor diet or lack of physical activity are inadequate to account for this dramatic and literal growth in our population, demonstrating the urgent need for new insight into mechanisms that may lead to obesity and altered energy homeostasis. Although it is well established that genetic variation plays a substantial role in the determination of body size and adiposity, the contribution of environmental alterations that have taken place in the last 20-30 years cannot be ignored. In addition to an increase in the abundance of low-quality, highly palatable, energy dense food, combined with a decrease in opportunities for exercise and physical activity, the average time spent sleeping is also decreasing. Recently, a number of studies have supported a link between the altered sleep/wake/eating patterns associated with our “24 hour” lifestyle and obesity.¹ According to the 2008 Annual Report from the National Sleep Foundation, obese individuals on average spent less time in bed (6.8 vs. 7.2 hours) and were more likely to sleep less than 6 hours per night (19% vs. 12%) compared to non-obese individuals. At present, the mechanisms by which altered daily rhythms are translated into increased adiposity are unknown.

Biological rhythms, such as sleep/wake cycles, are an integral component of virtually all aspects of life. These rhythms are controlled in large part by circadian clocks, intrinsically maintained molecular mechanisms that serve to condition the organism to changes in its environment. Though many studies have investigated the function of the central clock in the regulation of circadian physiology, our understanding of the role of peripheral clocks in whole body energy metabolism is just beginning to emerge. Recent studies have provided evidence that both central and peripheral circadian clocks likely regulate many physiologic functions, including insulin sensitivity, endocrine regulation, energy homeostasis, satiety signaling, and cellular proliferation.²⁻⁵ The phenotypic outcomes of circadian clock disruption are varied, with sleep deprivation and shift work associated with insulin resistance, increased hunger, alterations in adipocytokines, and increased risk for obesity, cardiovascular disease, and type 2 diabetes.⁶⁻⁸

Disruption of the circadian clock can result from alterations in the DNA sequence of clock component or clock-regulated genes and/or loss of synchrony between natural patterns of light/dark, sleeping, and eating. In animals, disruption of clock-component genes has been associated with altered feeding patterns, hormonal abnormalities, alterations in blood lipids and adipokines, alterations in lean and fat mass, defective glucose homeostasis, and reduced life span.^{4,5,9-11} Limited studies to date have investigated variation in the clock-component or clock-regulated genes for association to body fat and obesity in humans. Multiple variants in CLOCK have been reported to be associated with the development of the metabolic syndrome, type 2 diabetes and cardiovascular disease.¹²⁻¹⁴ DNA sequence variants in CLOCK, CK1E, GSK3B, PER2, and PER3 have been associated with sleep and mood disorders, bipolar disease, seasonal affective disorder, advanced sleep phase syndrome, depression, and morning-evening preference.¹⁵⁻¹⁹ While obesity has been associated with a number of these conditions, limited studies specifically investigating the role of variation within circadian clock genes in human obesity have been reported to date. The combination of genetic alterations in the circadian clock and behavioral dyssynchrony with the 24-h environment may have profound effects on overall health. This presentation will provide an overview of the current knowledge of the role of clock-associated genetic mutation/variation in the development of obesity and related co-morbidities.

References:

1. Vorona, RD, Winn, MP, Babineau, TW, Eng, BP, Feldman, HR & Ware, JC. (2005) Overweight and obese patients in a primary care population report less sleep than patients with a normal body mass index. *Arch Intern Med* 165, 25-30.
2. la Fleur, SE, Kalsbeek, A, Wortel, J, Fekkes, ML & Buijs, RM. (2001) A daily rhythm in glucose tolerance: a role for the suprachiasmatic nucleus. *Diabetes* 50, 1237-1243.

3. Cailotto, C, La Fleur, SE, Van Heijningen, C, Wortel, J, Kalsbeek, A, Feenstra, M et al. (2005) The suprachiasmatic nucleus controls the daily variation of plasma glucose via the autonomic output to the liver: are the clock genes involved? *Eur J Neurosci* 22, 2531-40.
4. Shimba, S, Ishii, N, Ohta, Y, Ohno, T, Watabe, Y, Hayashi, M et al. (2005) Brain and muscle Arnt-like protein-1 (BMAL1), a component of the molecular clock, regulates adipogenesis. *Proc Natl Acad Sci U S A* 102, 12071-6.
5. Turek, FW, Joshu, C, Kohsaka, A, Lin, E, Ivanova, G, McDearmon, E et al. (2005) Obesity and metabolic syndrome in circadian Clock mutant mice. *Science* 308, 1043-5.
6. Spiegel, K, Leproult, R & Van Cauter, E. (1999) Impact of sleep debt on metabolic and endocrine function. *Lancet* 354, 1435-9.
7. Vgontzas, AN, Papanicolaou, DA, Bixler, EO, Kales, A, Tyson, K & Chrousos, GP. (1997) Elevation of plasma cytokines in disorders of excessive daytime sleepiness: role of sleep disturbance and obesity. *J Clin Endocrinol Metab* 82, 1313-6.
8. Karlsson, BH, Knutsson, AK, Lindahl, BO & Alfredsson, LS. (2003) Metabolic disturbances in male workers with rotating three-shift work. Results of the WOLF study. *Int Arch Occup Environ Health* 76, 424-30.
9. Bunger, MK, Wilsbacher, LD, Moran, SM, Clendenen, C, Radcliffe, LA, Hogenesch, JB et al. (2000) Mop3 is an essential component of the master circadian pacemaker in mammals. *Cell* 103, 1009-17.
10. Kondratov, RV, Shamanna, RK, Kondratova, AA, Gorbacheva, VY & Antoch, MP. (2006) Dual role of the CLOCK/BMAL1 circadian complex in transcriptional regulation. *Faseb J* 20, 530-2.
11. Dallmann, R, Touma, C, Palme, R, Albrecht, U & Steinlechner, S. (2006) Impaired daily glucocorticoid rhythm in *Per1* (*Brd*) mice. *J Comp Physiol A Neuroethol Sens Neural Behav Physiol* 192, 769-75.
12. Sookoian, S, Gemma, C, Gianotti, TF, Burgueno, A, Castano, G & Pirola, CJ. (2008) Genetic variants of Clock transcription factor are associated with individual susceptibility to obesity. *Am J Clin Nutr* 87, 1606-15.
13. Monteleone, P, Tortorella, A, Docimo, L, Maldonato, MN, Canestrelli, B, De Luca, L et al. (2008) Investigation of 3111T/C polymorphism of the CLOCK gene in obese individuals with or without binge eating disorder: association with higher body mass index. *Neurosci Lett* 435, 30-3.
14. Scott, EM, Carter, AM & Grant, PJ. (2008) Association between polymorphisms in the Clock gene, obesity and the metabolic syndrome in man. *Int J Obes (Lond)* 32, 658-62.
15. Mishima, K, Tozawa, T, Satoh, K, Saitoh, H & Mishima, Y. (2005) The 3111T/C polymorphism of hClock is associated with evening preference and delayed sleep timing in a Japanese population sample. *Am J Med Genet B Neuropsychiatr Genet* 133B, 101-4.
16. Benedetti, F, Bernasconi, A, Lorenzi, C, Pontiggia, A, Serretti, A, Colombo, C et al. (2004) A single nucleotide polymorphism in glycogen synthase kinase 3-beta promoter gene influences onset of illness in patients affected by bipolar disorder. *Neurosci Lett* 355, 37-40.
17. Archer, SN, Robilliard, DL, Skene, DJ, Smits, M, Williams, A, Arendt, J et al. (2003) A length polymorphism in the circadian clock gene *Per3* is linked to delayed sleep phase syndrome and extreme diurnal preference. *Sleep* 26, 413-5.
18. Johansson, C, Willeit, M, Smedh, C, Ekholm, J, Paunio, T, Kieseppa, T et al. (2003) Circadian clock-related polymorphisms in seasonal affective disorder and their relevance to diurnal preference. *Neuropsychopharmacology* 28, 734-9.

19. Robilliard, DL, Archer, SN, Arendt, J, Lockley, SW, Hack, LM, English, J et al. (2002) The 3111 Clock gene polymorphism is not associated with sleep and circadian rhythmicity in phenotypically characterized human subjects. *J Sleep Res* 11, 305-12.

Maternal Imprinting and Metabolism: A Circadian Relationship?

Greg Sutton, PhD

Pennington Biomedical Research Center, LSU System, Baton Rouge, LA

The mechanisms linking intrauterine growth retardation (IUGR) with adulthood obesity and diabetes are unknown. These studies investigated energy metabolism in 8 wk and 5 month old male and female mice subjected to protein malnutrition in utero. We investigated the metabolic phenotype of IUGR offspring prior to the onset of obesity. We also investigated whether B6 mice subjected to prenatal protein deficiency also develop signs of Type II Diabetes. Undernourished pups from pregnant C57BL/6J dams fed a protein deficient diet (6 % protein, UO) were cross-fostered to lactating dams fed normal breeder chow. UO exhibited lower birth weight ($\approx 60\%$ of normal weight), but displayed rapid catch-up growth. 8 wk old chow-fed UO mice exhibit improved glucose clearance relative to CO. However, UO exhibited marked abnormalities in circadian rhythms of wheel running, feeding behavior and metabolic activity. Specifically, food intake, energy expenditure and the respiratory exchange ratio (RER) were increased in UO during the lights-on period. Expression of genes involved in hepatic lipid and glucose metabolism revealed that expression of Rev-erb α in liver was dramatically reduced in UO at 2 mo of age. Rev-erb α repressed genes involved in circadian regulation (Bmal1, Per2) and inflammation (plasminogen-activator inhibitor-1) were increased in UO mice. The same profile of gene expression was observed in male UO at 5 mo of age. UO mice exhibit a metabolic disorder involving abnormal circadian patterns of feeding behavior, increased lipogenesis and inflammation prior to obesity. Loss of Rev-erb α expression and function may be a key factor in metabolic dysregulation associated with IUGR.

Food Entrainment and the Circadian Regulatory Apparatus in Peripheral Tissues

Jeffrey M. Gimble¹, Z. Elizabeth Floyd¹, Sanjin Zvonic^{1*}, Xiyang Wu¹, Gang Yu¹, Brian C. Goh¹, Teddi Hebert^{1**}, Katie Hamel¹, Bruce A. Bunnell², Hui Xie¹, Steven R. Smith¹, Andrey A. Ptitsyn³

¹Pennington Biomedical Research Center, LSU System, Baton Rouge, LA; ²Tulane University, New Orleans, LA; ³Colorado State University, Ft. Collins, CO

*Currently at Tulane University, New Orleans, LA; **Currently at Louisiana State University Health Science Center-Shreveport, Shreveport, LA

Classical studies of human obesity and adipogenesis have focused on the transcriptional regulators C/EBP α and PPAR γ 2 and their downstream target genes relating to glucose and lipid metabolism. There is a growing awareness that circadian regulatory mechanisms play a contributory role in adipose tissue function. Studies in murine models have demonstrated the oscillatory expression of mRNAs encoding core circadian regulatory proteins (CCRP) in adipose depots [Ando et al., 2005; Bray and Young, 2007; Ptitsyn AA, 2006; Zvonic et al., 2006]. The adipose expression profile of the CCRP mRNAs can be significantly phase shifted by temporal restriction of food access [Goh BC and Hill MR, 2007]. In some obesity models, the CCRP oscillatory expression profile in adipose tissue is attenuated [Ando et al., 2005]. The same CCRP mRNAs were detected in subcutaneous adipose tissue of human subjects (n = 151). Analysis of the female Caucasian subjects (n = 116) found that expression of DBP, E4BP4, Rev-erb α , and PGC1 β all showed a trend (p < 0.1) or significant (p < 0.05) correlation with BMI in lean or young overweight/obese subjects; however, in older overweight/obese subjects, no correlation between CCRP mRNAs and BMI was detected. Consistent with these findings, human adipose derived stem cells (ASCs), capable of adipogenic, chondrogenic, and osteogenic differentiation, displayed a synchronized expression of CCRP mRNAs following exposure to dexamethasone or serum shock [Wu et al., 2007]. Similar observations were obtained in studies using bone marrow derived mesenchymal stem cells (BMSCs) from humans or mice [Wu et al., 2008]. Together, these studies indicate that human adipose tissue expresses the CCRP mRNAs in an oscillatory manner similar to that documented in murine models.

As observed in other peripheral, metabolically active tissues, at least 20% of the expressed mRNAs in subcutaneous adipose tissue exhibit an oscillatory expression profile [Ptitsyn AA, 2006; Zvonic et al., 2006]. A comparison of oscillatory mRNA in the transcriptome of four tissues (brown adipose, calvarial bone, liver, subcutaneous white adipose) revealed a shared set of ~180 transcripts common to each [Zvonic et al., 2007]. These included all members of the CCRP displayed on the microarray as well as mRNAs encoding critical rate limiting enzymes in glucose, iron, and lipid metabolism. An additional transcript in this shared subgroup displayed the intriguing name of "delta sleep inducing peptide immunoreactant (DSIPI)". Also known as "glucocorticoid induced leucine zipper (GILZ)", DSIPI was a protein first identified by antibodies directed against the nine amino acid delta sleep inducing peptide. The delta sleep inducing peptide was first discovered in the serum of sleeping mammals and was found to cause sleep when infused at high concentrations into otherwise awake animals. Pennington research Abba Kastin MD performed some of the pioneering work on delta sleep inducing peptide and demonstrated that its serum levels exhibited a robust circadian oscillation over a quarter century ago [Fischman et al., 1984; Graf and Kastin, 1984]; however, the protein source of the peptide remains unknown. Studies by Shi and his colleagues at the Medical College of Georgia have documented a regulatory role of DSIPI (GILZ) in the reciprocal regulation of adipogenesis and osteogenesis in BMSCs [Shi et al., 2003; Zhang et al., 2008]. In light of the growing interest in the relation between obesity risk and sleep, both DSIP and GILZ offer promising targets for future investigations in both pre-clinical and clinical models.

References:

Ando H, Yanagihara H, Hayashi Y, Obi Y, Tsuruoka S, Takamura T, Kaneko S, Fujimura A. 2005. Rhythmic mRNA Expression of Clock Genes and Adipocytokines in Mouse Visceral Adipose Tissue. *Endocrinology*.

Bray MS, Young ME. 2007. Circadian rhythms in the development of obesity: potential role for the circadian clock within the adipocyte. *Obes Rev* 8:169-81.

Fischman AJ, Kastin AJ, Graf MV. 1984. Circadian variation of DSIP-like material in rat plasma. *Life Sci* 35:2079-84.

Goh BC XWX, Ann E. Evans AE, Meagan L. Johnson ML, Molly R., Hill MR GJ. 2007. Food Entrainment of Circadian Gene Expression Altered in PPAR α -/- Brown Fat and Heart. *. Biochemical and Biophysical Research Communications* 360:828-833.

Graf MV, Kastin AJ. 1984. Delta sleep-inducing peptide (DSIP)-like material exists in peripheral organs of rats in large dissociable forms. *Proc Soc Exp Biol Med* 177:197-204.

Ptitsyn AA ZS, Conrad SA, Scott LK, Mynatt ML, Gimble JM. 2006. Circadian Clocks are Resounding in Peripheral Tissues. *PLoS Computational Biology* 2:e16.

Shi X, Shi W, Li Q, Song B, Wan M, Bai S, Cao X. 2003. A glucocorticoid-induced leucine-zipper protein, GILZ, inhibits adipogenesis of mesenchymal cells. *EMBO Rep* 4:374-80.

Wu X, Yu G, Parks H, Hebert T, Goh BC, Dietrich MA, Pelled G, Izadpanah R, Gazit D, Bunnell BA, Gimble JM. 2008. Circadian mechanisms in murine and human bone marrow mesenchymal stem cells following dexamethasone exposure. *Bone* 42:861-70.

Wu X, Zvonic S, Floyd ZE, Kilroy G, Goh BC, Hernandez TL, Eckel RH, Mynatt RL, Gimble JM. 2007. Induction of circadian gene expression in human subcutaneous adipose-derived stem cells. *Obesity (Silver Spring)* 15:2560-70.

Zhang W, Yang N, Shi XM. 2008. Regulation of mesenchymal stem cell osteogenic differentiation by glucocorticoid-induced leucine zipper (GILZ). *J Biol Chem* 283:4723-9.

Zvonic S, Ptitsyn AA, Conrad SA, Scott LK, Floyd ZE, Kilroy G, Wu X, Goh BC, Mynatt RL, Gimble JM. 2006. Characterization of peripheral circadian clocks in adipose tissues. *Diabetes* 55:962-70.

Zvonic S, Ptitsyn AA, Kilroy G, Wu X, Conrad SA, Scott LK, Guilak F, Pelled G, Gazit D, Gimble JM. 2007. Circadian Oscillation of Gene Expression in Murine Calvarial Bone. *J Bone Miner Res* 22:357-365.

The Circadian Biology of Cardiac Tissues: A Metabolic Model for Understanding Human Metabolic Syndrome

Martin E. Young, PhD

USDA/ARS Children's Nutrition Research Center, Baylor College of Medicine, Houston, TX

Diurnal variations have been firmly established in both cardiovascular physiology (e.g. heart rate, cardiac output) and pathophysiology (e.g. arrhythmias). As with myocardial contractile function, myocardial metabolism exhibits a marked diurnal variation. These phenomena have been attributed primarily to diurnal variations in extracardiac influences, including various neurohumoral factors, such as sympathetic and autonomic stimulation. Virtually every mammalian cell possesses an intrinsic circadian clock, a transcriptionally-based molecular mechanism capable of regulating multiple cellular functions. Circadian clocks allow individual cells to perceive the time of day. In doing so, this cell autonomous molecular mechanism confers the selective advantage of anticipation, enabling both rapid and appropriate responses to environmental stimuli upon their onset.

CLOCK mutant mice, which exhibit altered diurnal variations at multiple levels (including feeding-fasting cycles, neurohumoral factors), have been reported to develop multiple features of the Cardiometabolic Syndrome, suggesting that circadian clocks play critical roles in energy homeostasis. We have recently generated a cardiomyocyte-specific circadian clock mutant (CCM) mouse, as a means to identify the influences that this molecular mechanism has on myocardial physiology and pathophysiology. This model has exposed the cardiomyocyte circadian clock as a critical modulator in the responsiveness of the heart to fatty acids, at transcriptional, metabolic, and contractile function levels. We recently reported that the rodent heart exhibits increased susceptibility to fatty acid-induced contractile dysfunction during the less active/sleep (light) phase, a phenomenon that may be due to increased channeling of fatty acids into non-oxidative 'lipotoxic' pathways at this time. In contrast, exposure of the rodent heart to fatty acids during the more active/awake (dark) phase is associated with promotion of fatty acid oxidation, via transcriptional mechanisms. Through the use of CCM mice, studies within our laboratory suggest that both diurnal variations in fatty acid channeling, and in the transcriptional responsiveness of the heart to fatty acids, are mediated, at least in part, by the circadian clock within the cardiomyocyte. For example, the cardiomyocyte circadian clock directly regulates myocardial triglyceride metabolism, as evidenced by attenuated triglyceride syntheses for CCM hearts both *in vivo* (fasting-induced) and *ex vivo* (working mouse heart perfusions), through regulation of multiple triglyceride metabolism genes (e.g. *dgat2*, *adpn*).

Armed with the aforementioned knowledge, we have initiated studies designed to investigate whether restricting dietary lipid to distinct times of the day influences multiple parameters of the Cardiometabolic Syndrome. These studies suggest that restricting dietary lipid to the initiation of the more active/awake (dark) phase provides significant benefit at the levels of body weight gain/adiposity, glucose tolerance, skeletal muscle insulin sensitivity, and myocardial contractile function.

Taken together, our studies show that the circadian clock within the cardiomyocyte allows the heart to anticipate environmental stimuli (such as fatty acids), and that synchronization between the myocardium and its environment is enhanced by distinct meal feeding regimes. Impairment of the cardiomyocyte circadian clock (as observed during diabetes mellitus, ischemic heart disease) results in a loss of synchronization between the heart and its environment, potentially predisposing the myocardium to metabolic maladaptation and subsequent contractile dysfunction. Similarly, dyssynchronization of dietary lipid intake and circadian clock-mediated peripheral tissue fatty acid oxidation promotes Cardiometabolic Syndrome development.

Long-term Effects of Nightly Sleep Duration on Cognitive Performance, Alertness, Metabolism, and Obesity

Tracy L. Rupp, Ph.D. and Thomas J. Balkin, Ph.D.

Walter Reed Army Institute of Research, Department of Behavioral Biology, Silver Spring, MD

American adults report sleeping an average of 6.8 hours on weeknights¹—considerably less than the 8 h of sleep thought to be necessary to restore and sustain optimal daytime alertness. Whereas total and partial sleep loss each produce similar effects on objectively measured alertness and performance, the time course for recovery from chronic partial sleep loss is relatively extended.² Evidence also suggests that chronic sleep restriction leads to physiological changes that may be linked to obesity and diabetes.^{3,4,5,6} According to a review by Knutson and colleagues,⁷ the relationship between sleep loss, weight gain, and diabetes risk involves multiple pathways; 1) alterations in glucose, 2) upregulation of appetite, and 3) decreased energy expenditure.

Here we present evidence that the extent to which sleep restriction impairs objectively measured alertness and performance, and the rate at which these impairments are subsequently reversed by recovery sleep, varies as a function of the amount of nightly sleep obtained prior to the sleep restriction period.⁸ We then discuss implications of the present findings for understanding the potential effects of sleep extension and partial sleep loss on metabolism and obesity; including that prior sleep extension might mediate (and potentially buffer) the multiple pathways connecting sleep loss to weight gain and diabetes risk, specifically with regards to alterations in glucose and the upregulation of appetite.

References:

1. National Sleep Foundation. Sleep in America Poll. 2005. Washington DC. See <http://sleepfoundation.org>.
2. Belenky G, Wesensten NJ, Thorne DR, et al. Patterns of performance degradation and restoration during sleep restriction and subsequent recovery: a sleep dose-response study. *J Sleep Res* 2003; 12:1-12.
3. Spiegel K, Leproult M, Van Cauter E. Impact of sleep debt on metabolic and endocrine function. *Lancet* 1999; 354:1435-1439.
4. Spiegel K, Knutson K, Leproult R, et al. Sleep loss: a novel risk factor for insulin resistance and Type 2 diabetes. *J Appl Physiol* 2005; 99:2008-2019.
5. Chaput JP, Despres JP, Bouchard C, et al. The association between sleep duration and weight gain in adults: A 6-year prospective study from the Quebec family study. *Sleep* 2008; 31:517-523.
6. Ayas NT, White DP, Al-Delaimy WK, et al. A prospective study of self-reported sleep duration and incident diabetes in women. *Diabetes Care* 2004; 27:282-284.
7. Knutson KL, Spiegel K, Penev P, Van Cauter E. The metabolic consequences of sleep deprivation. *Sleep Medicine Reviews* 2007; 11:163-178.
8. Rupp TL, Wesensten NJ, Bliese PD, Balkin TJ. Banking sleep: Realization of benefit during subsequent sleep restriction and recovery. *Sleep* 2009; 32:311-21.

Obesity and Sleep Disturbances

Alexandros Vgontzas, MD
Penn State College of Medicine, Hershey, PA

Obesity, excessive daytime sleepiness (EDS), and self-reported short sleep duration appear to be on the rise, while there is evidence that obesity and these sleep disorders are strongly connected. In our presentation, we will review data that challenge the common belief that the sleep apnea and sleep loss, frequently associated with obesity, are the primary determinants of obesity-related objective daytime sleepiness and subjective fatigue (tiredness without increased sleep propensity). Specifically, obesity is associated with objective and subjective EDS regardless of the presence of sleep apnea. The association between obesity and EDS was confirmed in recent studies of large random samples of the general population or clinical samples, which showed that the primary determinants of subjective EDS were depression, metabolic disturbances, i.e., obesity/diabetes and insulin resistance, and lack of physical activity, and, secondarily, sleep apnea or sleep loss. Paradoxically, within the obese, with or without sleep apnea, those who slept objectively better at night are sleepier (objectively) during the day than those who slept worse. The distinguishing factor between those that slept better vs. those that slept worse appears to be level of emotional stress. Furthermore, many studies reported that obesity is associated with self-reported short sleep duration; however, it appears that short sleep duration is a marker of emotional stress rather than a reflection of true sleep loss. Based on these data, we propose that obesity-related deeper sleep and objective EDS are primarily related to metabolic disturbances, whereas obesity-related poorer sleep and subjective fatigue appear to be the result of psychological distress.

Furthermore, based on data from studies in normal controls and patients with sleep disorders, it appears that the interaction of the hypothalamic-pituitary-adrenal (HPA) axis and pro-inflammatory cytokines determines the level of sleep/arousal within the 24-hour cycle, i.e., "eucortisolemia" or "hypocortisolemia" plus hypercytokinemia is associated with high sleep efficiency and objective sleepiness, whereas "hypercortisolemia" plus hypercytokinemia is associated with low sleep efficiency and fatigue.

In conclusion, we propose that the above-reviewed data provide the basis for a meaningful phenotypic and pathophysiologic subtyping of obesity. One subtype is associated with emotional distress, poor sleep, fatigue, HPA axis "hyperactivity," and hypercytokinemia while the other is associated with nondistress, better sleep but more sleepiness, HPA axis "normo or hypoactivity," and hypercytokinemia. This proposed subtyping may lead to novel, preventive and therapeutic strategies for obesity and its associated sleep disturbances.

References:

- Bixler EO, Vgontzas AN, Lin HM, Calhoun SL, Vela-Bueno A, Kales A. 2005. Excessive daytime sleepiness in a general population sample: the role of sleep apnea, age, obesity, diabetes, and depression. *J Clin Endocrinol Metab* 90(8):4510-5.
- Guan Z, Vgontzas AN, Bixler EO, Fang J. 2008. Sleep is increased by weight gain and decreased by weight loss in mice. *Sleep* 31(5):627-33.
- Lauderdale DS, Knutson KL, Yan LL, Rathouz PJ, Hulley SB, Sidney S, Liu K. 2006. Objectively measured sleep characteristics among early-middle-aged adults: the CARDIA study. *Am J Epidemiol* 164(1):5-16.
- Mårin P, Darin N, Amemiya T, Andersson B, Jern S, Björntorp P. 1992. Cortisol secretion in relation to body fat distribution in obese premenopausal women. *Metabolism* 41(8):882-6.
- Punjabi NM, O'hearn DJ, Neubauer DN, Nieto FJ, Schwartz AR, Smith PL, Bandeen-Roche K. 1999. Modeling hypersomnolence in sleep-disordered breathing. A novel approach using survival analysis. *Amer J Respir Crit Care Med* 159(6):1703-9.
- Resta O, Foschino-Barbaro MP, Legari G, Talamo S, Bonfitto P, Palumbo A, Minenna A, Giorgino R, De Pergola G. 2001. Sleep-related breathing disorders, loud snoring and excessive daytime sleepiness in obese subjects. *Int J Obes Relat Metab Disord* 25(5):669-75.

Spiegel K, Tasali E, Penev P, Van Cauter E. 2004. Brief communication: Sleep curtailment in healthy young men is associated with decreased leptin levels, elevated ghrelin levels, and increased hunger and appetite. *Ann Intern Med* 141(11):846-50.

Stunkard AJ, Faith MS, Allison KC. 2003. Depression and obesity. *Biol Psychiatry* 54(3):330-7.

Theorell-Haglöw J, Lindberg E, Janson C. 2006. What are the important risk factors for daytime sleepiness and fatigue in women? *Sleep* 29(6):751-7.

Vgontzas AN, Tan TL, Bixler EO, Martin LF, Shubert D, Kales A. 1994. Sleep apnea and sleep disruption in obese patients. *Arch Intern Med* 154(15):1705-11.

Vgontzas AN, Papanicolaou DA, Bixler EO, Kales A, Tyson K, Chrousos GP. 1997. Elevation of plasma cytokines in disorders of excessive daytime sleepiness: role of sleep disturbance and obesity. *J Clin Endocrinol Metab* 82(5):1313-6.

Discussion Questions:

1. Although the impact of clock gene disruption on metabolism and related health problems has been highlighted by a number of studies, a recurring question is whether the observed defects are due to clock dysfunction (e.g. via deregulated clock-controlled genes) or to clock-independent roles of these so-called clock genes. The same question applies to all other health issues occurring upon clock gene dysfunction.
2. Does sleep extension a) protect against the effects of subsequent sleep loss on metabolism and/or b) help reverse metabolic effects associated with prior sleep debt?
3. Are there age-related changes in the effects of chronic sleep on metabolism and obesity?
4. What specific countermeasures (pharmacological or behavioral) might protect against the metabolic effects of chronic insufficient sleep?
5. How might laboratory versus field studies best capture the long-term consequences of chronic insufficient sleep on metabolism?
6. Are the metabolic phenotypes observed in the various clock-deficient mice the result of loss of the clock *per se*, or due to other pleiotropic effects of the clock genes (or a combination of the two)?
7. What consideration are people giving to background strain when conducting metabolic studies in circadian transgenics?
8. What is the evidence that the night eating syndrome involves a disturbance of the glucocorticoid system, the melatonergic system and the serotonergic system?
9. What do you think is the reason for the lack of knowledge on the part of physicians of this easily diagnosed and readily treated disorder?

CONTACT INFORMATION

CO-CHAIRS:

Molly Bray, PhD

Associate Professor of Pediatrics
Baylor College of Medicine
Children's Nutrition Research Center
1100 Bates, Ste. 8070
Houston, TX 77030
Telephone: 713-798-9310
Email: mbray@bcm.tmc.edu

Jeffrey Gimble, MD, PhD

Professor, Stem Cell Biology
Pennington Biomedical Research Center
Louisiana State University System
6400 Perkins Rd
Baton Rouge, LA 70808
Telephone: 225-763-3171
Email: gimblejm@pbrc.edu

Andrew Young, PhD

Chief, Military Nutrition Division
U.S. Army Research Institute of Environmental Medicine
Natick, MA 01760
Telephone: 508-233-5141
Email: Andrew.Young@amedd.army.mil

Speakers and Invited Scientific Colleagues

Thomas Balkin, PhD

Chief, Department of Behavioral Biology
Walter Reed Army Institute of Research
503 Robert Grant Ave., Room # 2A26
Silver Spring, MD 20910
Telephone: 301-319-9350
Email: Thomas.Balkin@amedd.army.mil

Joseph Bass, MD, PhD

Assistant Professor, Medicine and Neurobiology and Physiology
Northwestern University Feinberg School of Medicine
2205 Tech Drive, Hogan 2-100
Evanston, IL 60208
Telephone: 847-491-5571
Email: j-bass@northwestern.edu

Julia Boyle, PhD

Director of Surrey Clinical Research Center, Health & Medical Sciences
University of Surrey
Guildford Surrey GU2 7XP
Telephone: 44 0 1483 689783
Email: J.Boyle@surrey.ac.uk

Andrew Butler, PhD

Associate Professor, Department of Metabolism and Aging
The Scripps Research Institute
130 Scripps Way
Jupiter, FL 33458
Telephone: 561-228-2957
Email: abutler@scripps.edu

Leticia Castillo, MD

Associate Professor, Pediatrics
Baylor College of Medicine
1100 Bates
Houston, TX 77030
Telephone: 832-826-6225
Email: lcastill@bcm.edu

Nicolas Cermakian, PhD

Professor, Douglas Mental Health University Institute
Department of Psychiatry, McGill University
6875, LaSalle Blvd
Montreal, Canada H4H 1R3
Telephone: 1-514-761-6131 #4936
Email: nicolas.cermakian@mcgill.ca

Karyn Esser, PhD

Associate Professor, Department of Physiology
University of Kentucky
Lexington, KY 40536
Telephone: 859-323-8107
Email: karyn.esser@uky.edu

Jidong Fang, MD, PhD

Associate Professor of Psychiatry, Milton S. Hershey Medical Center
Penn State
500 University Drive
Hershey, PA 17033
Telephone: 717-531-5435
Email: jfang@psu.edu

James Gangwisch, PhD

Assistant Professor, Department of Psychiatry, NYSPI
Columbia University
1051 Riverside Drive
Room 2304, Mail Code 74
New York, NY 10032
Telephone: 212-543-6552
Email: jeg64@columbia.edu

Carla Green, PhD

Professor, Department of Biology
University of Virginia
Room 275, Gilmer Hall
PO Box 400328
Charlottesville, VA 22904
Telephone: 434-982-5436
Email: cbg8b@virginia.edu

Rich Heyman

Research Scientist and Consultant
2825 Crystal Ridge Rd
Encinitas, CA 92024
Telephone: 858-735-4581
Email: rich.heyman@gmail.com

Jonathan Johnston, PhD

Faculty, Health and Sciences
The University of Surrey
Guilford, Surrey
GU2 7XH United Kingdom
Telephone: +44 (0) 1483 - 686470
Email: j.johnston@surrey.ac.uk

Chintan Kikani, PhD

Postdoctoral Fellow
University of Utah School of Medicine
15N Medical Drive East, Room #4208
Salt Lake City, UT 84109
Telephone: 801-581-5552
Email: chintan.kikani@utah.edu

Biliana Marcheava

Graduate Student
Northwestern University
2200 Campus Drive
Evanston, IL 60208
Telephone: 847-467-5973
Email: b-marcheva@northwestern.edu

Sanjay Patel, MD

Assistant Professor
Case Western Reserve University
11400 Euclid Ave. Room 290-D
Cleveland, OH 44106
Telephone: 216-844-6258
Email: sanjay.r.patel@case.edu

Ignacio Provencio, PhD

Associate Professor, Department of Biology
University of Virginia
281 Gilmer Hall
Charlottesville, VA 22904
Telephone: 434-924-4412
Email: ip7m@virginia.edu

Andrew Ptitsyn, PhD

Assistant Professor, Center for Bioinformatics
Colorado State University
Fort Collins, CO 80523-1682
Telephone: 970-491-0878
Email: ptitsyn@lamar.colostate.edu

Tracy Rupp, PhD

Research Psychologist, Department of Behavioral Biology
Walter Reed Army Institute of Research
Division of Psychiatry and Neuroscience
503 Robert Grant Avenue
Silver Spring, MD 20910
Telephone: 301-319-9352
Email: tracy.rupp@amedd.army.mil

Jared Rutter, PhD

Assistant Professor, Department of Biochemistry
University of Utah School of Medicine
15 N. Medical Drive East Room 4280
Salt Lake City, UT 84112-5650
Telephone: 801-581-3340
Email: rutter@biochem.utah.edu

Kai-Florian Storch, PhD

Assistant Professor, Douglas Mental Health Institute
McGill University
6875 LaSalle Blvd
Montreal, Quebec Canada H4H 1R3
Telephone: 514-761-6131#3940
Email: florian.storch@mcgill.ca

Albert Stunkard, MD

Professor of Psychiatry, Department of Psychiatry
Weight and Eating Disorders Program
University of Pennsylvania
3600 Market Street, Rm. 734
Philadelphia, PA 19104-2648
Telephone: 215-898-7314
Email: stunkard@mail.med.upenn.edu

Greg Sutton, PhD

Instructor, Protein Deficiency and Developmental Biology
Pennington Biomedical Research Center
6400 Perkins Rd.
Baton Rouge, LA 70808
Telephone: 225-763-3121
Email: greg.sutton@pbrc.edu

Joseph Takahashi, PhD

Professor, WCAS Neurobiology & Physiology
Northwestern University
2205 Tech Drive, 2-141, EV3520
Evanston, IL 60208
Telephone: 847-491-4598
Email: j-takahashi@northwestern.edu

Michael Twery, PhD

Director, National Center on Sleep Disorders Research
NHLBI/NIH/DHHS
6701 Rockledge Drive, MSC 7952
Bethesda, MD 20892-7952
Telephone: 301-435-0199
Email: twerym@nhlbi.nih.gov

Alexandros Vgontzas, MD

Director, Sleep Research & Treatment Center
Professor of Psychiatry, Endowed Chair in Sleep Disorders Medicine
Penn State College of Medicine
500 University Drive, H073
Hershey, PA 17033-0850
Telephone: 717 5318515
Email: avgontzas@psu.edu

Martin Young, PhD

Assistant Professor, Children's Nutrition Research Center
Baylor College of Medicine
One Baylor Plaza, Mail Stop BCM320
Houston, TX 77030
Telephone: 713-798-7567
Email: meyoung@bcm.tmc.edu